

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/82932 A2(51) International Patent Classification⁷: **A61K 31/661**,
31/6615, 31/662, 31/7084, 31/706, 31/7076

(21) International Application Number: PCT/US01/13931

(22) International Filing Date: 30 April 2001 (30.04.2001)


(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/200,843 1 May 2000 (01.05.2000) US
60/230,263 6 September 2000 (06.09.2000) US
60/233,025 15 September 2000 (15.09.2000) US(71) Applicant: **HEALTHPARTNERS RESEARCH FOUNDATION** [US/US]; 8100 34th Avenue South, Minneapolis, MN 55440-1524 (US).(72) Inventors: **FREY, William, H., II.**; 4800 Centerville Road, Apt. 216, White Bear Lake, MN 55127 (US).**FAWCETT, John, Randall**; 928 Grand Avenue, # 305, St. Paul, MN 55105 (US). **THORNE, Robert, Gary**; 2610 Fremont Avenue South #106, Minneapolis, MN 55408 (US). **CHEN, Xueqing**; 1166 Filfield Avenue, Apt. S2, St. Paul, MN 55108 (US).(74) Agent: **FORREST, Peter, R.**; Gray, Plant, Mooty, Mooty & Bennett, P.A., P.O. Box 2906, Minneapolis, MN 55402-2906 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: METHODS AND COMPOSITIONS FOR ENHANCING CELLULAR FUNCTION THROUGH PROTECTION OF TISSUE COMPONENTS

Pyrophosphate Protects the mAChR from
inactivation by the LMW Inhibitor
(n = 3)

(57) Abstract: Methods and compositions for enhancing cellular function through protection of tissue components, such as receptors, proteins, lipids, nucleic acids, carbohydrates, hormones, vitamins, and cofactors, by administering pyrophosphate analogs or related compounds.

WO 01/82932 A2

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5

METHODS AND COMPOSITIONS FOR ENHANCING CELLULAR FUNCTION THROUGH PROTECTION OF TISSUE COMPONENTS

10

Cross-Reference to Related Applications

This application claims the benefit of provisional application number 60/200,843 filed May 1, 2000; provisional application number 60/233,263 filed September 6, 2000; and provisional application number 60/233,025 filed September 15, 2000; each entitled "Methods and Compositions for Enhancing Cellular Function and Protecting Receptor."

15

Background of the Invention

Cellular function depends on the maintenance of intact cellular components including: receptors, proteins, lipids, nucleic acids, carbohydrates, hormones and cofactors. Cellular receptors, including cell surface receptors, mediate communication within and between cells, tissues and organs within a living system. Cellular receptors also provide a means to signal a living system, tissues, organs, cells, and subcellular compartments. Receptors are molecules or macromolecules that bind or interact with agents to alter or enhance their function. Many receptors are membrane bound proteins, which require not only that their protein structure be intact but also that the membrane lipids and carbohydrates be intact and functional. Through various signaling mechanisms, the messages sent by the receptor, either in the presence or absence of an interacting or bound agent, can be transmitted. Following receptor activation, signalling also requires intact cellular proteins, lipids, nucleic acids and carbohydrates in order for the message to be properly received.

Often as a result of damage, the ability of cellular receptors to interact with or bind various agents is decreased, resulting in an impairment of vital intrinsic and extrinsic communication. Damage to cellular receptors and other cellular components diminishes the ability of a receptor to bind agents and elicit a communication or signaling event. This can result in damage or death to cells, resulting in damage or diseases of tissues, organs and living systems. Accordingly, there is a need for a means to protect receptors and other cellular components from damage and to increase the efficacy of agents that exert their effects through cellular receptors.

10

Summary of the Invention

The invention provides methods for enhancing cellular function through protection of tissue components and/or increasing the efficacy of a therapeutic agent in a subject in need thereof. The method includes administering a composition, such as a pharmaceutical composition, of a pyrophosphate analog. In a second embodiment, the method includes administering a composition, such as a pharmaceutical composition, of a protective agent.

Preferably, the invention provides a method for protecting a muscarinic acetylcholine receptor (mAChR) and/or increasing the efficacy of an agent that directly or indirectly affects a mAChR in a subject in need thereof. Suitable agents that directly or indirectly affect a muscarinic receptor include anticholinesterase agents, muscarinic agonists, allosteric regulators of a muscarinic receptor, muscarinic antagonists, and neurotrophic and neuritogenic factors that are similar to naturally occurring nerve growth promoting substances. In one embodiment, the invention provides a method to protect a mAChR and/or increase the efficacy of agents that directly or indirectly affect a mAChR in the central nervous system (CNS) of a subject in need thereof. Preferably, a muscarinic

receptor is protected from an endogenous low molecular weight inhibitor from Alzheimer's brain tissue, a metal, or oxidative stress. In another embodiment, the invention provides a method to protect a mAChR and/or increase the efficacy of agents that directly or indirectly affect a mAChR not in the CNS of a subject in a need thereof.

- 5 In a first embodiment, the method includes administering a pyrophosphate analog. In a second embodiment, the method includes administering a protective agent.


The invention also provides a method for increasing the efficacy of a therapeutic agent, preferably a neurologic agent, in a subject in need thereof. In a first embodiment, this method includes administering a pyrophosphate analog. In this embodiment, the

10 increased efficacy of the neurologic agent preferably results from protection of a muscarinic receptor caused or induced by the pyrophosphate analog. In a second embodiment, this method includes administering a protective agent. In this second embodiment, the increased efficacy of the neurologic agent preferably results from protection of a muscarinic receptor caused or induced by the protective agent. In each

15 embodiment, the subject preferably is concurrently receiving, has recently received, or will soon receive a neurologic agent such as nerve growth factor (NGF), insulin growth factor (IGF-1), brain derived neurotrophic factor (BDNF), fibroblast growth factor (FGF), and the like; certain other known neurotrophins and neuroprotectants; and medications used for stroke, Alzheimer's disease, Parkinson's disease, ALS, traumatic


20 brain or spinal cord injury, cancer, diabetes, neuropathies, hypertension, irritable bowel syndrome; diseases or disorders of the heart and smooth muscles, blood, blood vessels, glands or bones and other disorders. Preferably, the therapeutic agent directly or indirectly affects a mAChR. Such agents include an anticholinesterase agent, a muscarinic agonist, and a muscarinic antagonist.

Pyrophosphate analogs that can be employed in the appropriate embodiment of the method of the invention include compounds of Formula I:



where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²), and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900. Compounds of Formula I in which R¹ is a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²); or R² is a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl can be referred to as substituted pyrophosphate analogs. Compounds of Formula I can also include substituted pyrophosphate analogs such as a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, and a pyrimidine acyclonucleoside. The compound of Formula I can be any pharmaceutically acceptable salt or basic addition salt. Preferably, X is O, CH₂, NH, or S; R¹ is H; R² is H; and n is 1-6. More preferably the pyrophosphate analog is pyrophosphate or imidodiphosphate.

Additional pyrophosphate analogs include compounds of Formula II:



where n=2-4; X is O, RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, such as CH₃ or (CH₂)_mNH₂ where m=1-6.

Further included are bisphosphonic acids, which are also known as bisphosphonates, where X is preferably RCR^1 , where R and R^1 groups are chosen independently from OH, $\text{H}_2\text{N}(\text{CH}_2)_2$, or CH_3 . For example, RCR^1 can be $\text{H}_2\text{N}(\text{CH}_2)_2\text{C}(\text{OH})$ or CH_3COH . More specifically, the bisphosphonates include etidronic acid ((1-

- 5 Hydroxyethylidene)bisphosphonic acid) and pamidronic acid ((3-Amino- 1-hydroxypropylidene)bisphosphonic acid) where preferably $n=2$.

Yet more additional pyrophosphate analogs include substituted pyrophosphate analogs such as an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, and an inositol hexaphosphate.

- 10 Suitable protective agents that can be employed in an embodiment of the method of the invention include a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavonoid, a combination thereof, or a pharmaceutically acceptable salt thereof; a heme binding compound, such as hemopexin, lipopexin, a lipoprotein, or ApoE-2; and a heme oxygenase, such as heme oxygenase-1 or heme oxygenase-2, biliverdin reductase, a
- 15 catalase, a peroxidase, a vector encoding a biliverdin reductase, a vector encoding a heme oxygenase (e.g. a vector encoding a heme oxygenase-1 or a vector encoding a heme oxygenase-2), a vector encoding a catalase, a vector encoding a peroxidase, or a combination thereof Biliverdin reductase can be administered alone or in combination with a heme oxygenase. Heme oxygenases include recombinant heme oxygenase.
- 20 Preferably, a heme oxygenase is a human heme oxygenase.

- The method of the invention can treat or prevent a CNS disorder. Preferably, the method of the invention can treat or prevent neurodegeneration, can improve memory and cognition, can treat or prevent brain deterioration or cognitive and memory loss associated with aging, or can treat or prevent Alzheimer's Disease, Parkinson's disease,
- 25 Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear

WO 01/02 1
palsy, amyotrophic lateral sclerosis, affective disorders, anxiety disorders, and/or
schizophrenia; nerve damage from cerebrovascular disorders such as stroke or
atherosclerosis in the brain or spinal cord, from CNS infections including meningitis and
HIV, from tumors of the brain and spinal cord, prion diseases, and CNS disorders
5 resulting from ordinary aging (e.g., anosmia), brain injury, or spinal cord injury.

In another embodiment, the method of the invention can treat or prevent a disease
or disorder not of the CNS. Preferably, the method of the invention can treat or prevent
cancer, or neuropathies or diseases or disorders of the heart, smooth muscles, blood,
blood vessels, glands, or bones. Such diseases or disorders include hypertension,
10 myocardial infarction, ischemic heart disease, congestive heart failure, cardiac
arrhythmias, cancer, irritable bowel syndrome, diverticular disease, urinary incontinence,
esophageal achalasia, chronic obstructive airways disease, xerostomia, diabetes mellitus,
Sjogren's syndrome or dry eye syndrome which involves decreased secretion of tears by,
for example, the lacrimal glands, Paget's disease, hereditary hematochromatosis or a
15 non-CNS disorder resulting from normal aging.

In another embodiment, the method of the invention treats infections, including
(without limitation) bacterial, fungal, alga, or algae infections. Such infections can occur
in plants (for which a preferred embodiment of the invention employs imidodiphosphate
as the pyrophosphate analog), animals, or mammals.

20

Brief Description of the Figures

Figure 1 illustrates protection of a mAChR by pyrophosphate. Pyrophosphate
protected the mAChR from inactivation by the endogenous low molecular weight
inhibitor. Pyrophosphate protected the receptor from loss of antagonist (³H-QNB
25 (quinulidiny benzilate)) binding.

Figure 2 illustrates protection of a mAChR by pyrophosphate. Pyrophosphate protected the mAChR from inactivation by heme and peroxide. Pyrophosphate protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 3 illustrates protection of a mAChR by pyrophosphate. Pyrophosphate
5 protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Pyrophosphate protected the receptor from loss of agonist (oxotremorine) binding.

Figure 4 illustrates protection of a mAChR by imidodiphosphate.
Imidodiphosphate protected the mAChR from inactivation by the endogenous low
10 molecular weight inhibitor. Imidodiphosphate protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 5 illustrates protection of a mAChR by guanylimidodiphosphate.
Guanylimidodiphosphate protected the mAChR from inactivation by the endogenous low
molecular weight inhibitor. Guanylimidodiphosphate protected the receptor from loss of
15 antagonist (^3H -QNB) binding.

Figure 6 illustrates protection of a mAChR by adenylylimidodiphosphate.
Adenylylimidodiphosphate protected the mAChR from inactivation by the endogenous
low molecular weight inhibitor. Adenylylimidodiphosphate protected the receptor from
loss of antagonist (^3H -QNB) binding.

20 Figure 7 illustrates protection of a mAChR by tripolyphosphate.
Tripolyphosphate protected the mAChR from inactivation by the endogenous low
molecular weight inhibitor. Tripolyphosphate protected the receptor from loss of
antagonist (^3H -QNB) binding.

Figure 8 illustrates protection of a mAChR by bilirubin. Bilirubin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Bilirubin protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 9 illustrates protection of a mAChR by bilirubin. Bilirubin protected the
5 mAChR from inactivation by heme and peroxide. Bilirubin protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 10 illustrates protection of a mAChR by bilirubin. Bilirubin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Bilirubin protected the receptor from loss of agonist (oxotremorine) binding.

10 Figure 11 illustrates protection of a mAChR by biliverdin. Biliverdin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Biliverdin protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 12 illustrates protection of a mAChR by carnosol, Carnosol protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Carnosol
15 protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 13 illustrates protection of a mAChR by quercetin. Quercetin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Quercetin protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 14 illustrates protection of a mAChR by myricetin. Myricetin protected
20 the mAChR from inactivation by the endogenous low molecular weight inhibitor. Myricetin protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 15 illustrates protection of a mAChR by catalase. Catalase protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Catalase protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 16 illustrates protection of a mAChR by catalase. Catalase protected the mAChR from inactivation by heme and peroxide. Catalase protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 17 illustrates protection of a mAChR by a peroxidase. The peroxidase
5 protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. The peroxidase protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 18 illustrates protection of a mAChR by a peroxidase. The peroxidase protected the mAChR from inactivation by heme and peroxide. The peroxidase protected
10 the receptor from loss of antagonist (^3H -QNB) binding.

Figure 19 illustrates protection of mAChR by pamidronate. Pamidronate protected the mAChR from inactivation by the endogenous low molecular weight inhibitor.

Figure 20 illustrates protection of a mAChR by pyrophosphate. Pyrophosphate
15 protected the mAChR from damage by the metal lead in the form of PbCl_2 . Pyrophosphate protected the receptor from loss of antagonist (^3H -QNB (quinulidiny benzilate)) binding.

20

Detailed Description of the Invention

Definitions

As used herein, "cholinesterase" refers to an enzyme capable of hydrolyzing acetylcholine and includes acetylcholinesterase.

As used herein, "agonist" refers to an agent that binds to or interacts with a
25 receptor and elicits a response transduced through the receptor. Agonist includes full

agonists, partial agonists, and inverse agonists. A full agonist is an agent that can elicit a maximal response from a receptor. A partial agonist is an agent that can elicit, at best, a less than maximal response from a receptor. An inverse agonist is an agent that produces a response that is opposite that of a full or partial agonist. For example, if agonist
5 binding or interaction with a receptor results in increased concentration of cAMP within a cell, then inverse agonist binding or interaction with the same receptor will result in a decreased concentration of cAMP within the cell.

As used herein, “antagonist” refers to an agent that is capable of partially or completely inhibiting, or reversing, the effect of an agonist on a receptor.

10 As used herein, “allosteric modifier” refers to an agent that binds or interacts with a site other than the agonist binding site of a receptor and modifies the ability of an agonist or an antagonist to elicit or inhibit, respectively, a response transduced through a receptor, without itself eliciting a response.

As used herein, “tissue component” includes receptors, proteins, lipids, nucleic
15 acids, carbohydrates, hormones, vitamins, and cofactors.

As used herein, “receptor” refers to any molecule or macromolecule within or on a cell that interacts with another molecule or macromolecule to confer a response or transduce a signal and includes nuclear receptors, mitochondrial receptors, cytoplasmic receptors, and cell surface receptors. Receptors include receptors for neurotrophins
20 (including, without limitation, nerve growth factor, neurotrophins 3, 4, and/or 5 (NT-3, NT-4 and/or NT-5) and brain derived growth factors); neurotransmitters; hormones; steroids; local mediators such as nitric oxide, carbon monoxide, histamine, and growth factors like insulin, insulin-like growth factor-I, fibroblast growth factors, ciliary neurotrophic factor, glia-derived neurotrophic factor, glia-derived nexin, cholinergic
25 enhancing factor, transforming growth factors, activity dependent neurotrophic factor,

neurokines, gangliosides, phosphatidylserine, PDGF (platelet derived growth factor) and EGF (epidermal growth factor); benzodiazepines; arachidonic acid; purines (including, without limitation, adenosine and ATP); nucleotides and cyclic nucleotides; calcium and other divalent cations; odorants; antisense oligonucleotides; opiates; cannabinoids; glutamate; melatonin; angiotensin II; secretin; vasoactive intestinal peptide; cholecystikinin; ACTH; vasopressin; thrombin; ion channels; and the like. Receptors also include but are not limited to G-protein-coupled receptors, ion-channel-linked receptors and enzyme-linked receptors.

As used herein, "protecting a receptor" refers to protecting the physical integrity of a receptor and/or the function of a receptor, such as enhancing the function of a receptor; or maintaining the ability of the receptor to respond to agonists, to respond to antagonists, to transmit a message to the interior of a cell, or to send a signal within a cell, cell nucleus, or mitochondria.

As used herein, "central nervous system" (CNS) refers to the brain and spinal cord and associated tissues.

As used herein, "disorders and diseases of the CNS" refers to brain diseases such as Alzheimer's disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, affective disorders, anxiety disorders, and/or schizophrenia; cell damage; nerve damage from cerebrovascular disorders such as stroke in the brain or spinal cord, from CNS infections including meningitis and HIV, from tumors of the brain and spinal cord, prion diseases, and CNS disorders resulting from ordinary aging (e.g., anosmia), brain injury, or spinal cord injury.

As used herein, a disease or disorder that relates to or is caused at least in part by dysfunction, alteration, or loss of one or more G-protein coupled receptors refers to

Alzheimer's disease; Parkinson's disease; drug addiction, such as opiate addiction or cannabinoid abuse; pain; Sjogren's or dry eye syndrome; heart diseases including congestive heart failure, myocardial infarction, cardiac arrhythmia; diseases of smooth muscle organs or glands such as irritable bowel syndrome, colitis, hypertension, erectile dysfunction, diabetes, obesity, blood coagulation disorders; and the like.

An "effective amount" of agent is an amount sufficient to prevent, treat, reduce and/or ameliorate the symptoms and/or underlying causes of any of the above disorders or diseases. In some instances, an "effective amount" is sufficient to eliminate the symptoms of those diseases and, perhaps, overcome the disease itself. Preferably, an effective amount of an agent yields a tissue concentration in the range of about 10^{-7} molar to about 10^{-5} molar, but the concentrations may be greater provided that toxicity is avoided.

In the context of the present invention, the terms "treat" and "therapy" and the like refer to alleviate, slow the progression, prophylaxis, attenuation or cure of existing disease. Prevent, as used herein, refers to putting off, delaying, slowing, inhibiting, or otherwise stopping, reducing or ameliorating the onset of such diseases or disorders. It is preferred that a large enough quantity of the agent be applied in non-toxic levels in order to provide an effective level of activity against the disease. The method of the present invention may be used with any animal, such as a mammal or a bird (avian), more preferably a mammal. Poultry are a preferred bird. Exemplary mammals include, but are not limited to rats, cats, dogs, horses, cows, sheep, pigs, and more preferably humans.

Protecting a Tissue Component

The invention provides a method for protecting any biomolecule or tissue component including a protein, a lipid, a nucleic acid, a carbohydrate, a hormone and the

like. The invention is best illustrated, but is not limited to, the example of the protection of a receptor, preferably a muscarinic receptor, preferably a mAChR. In a first embodiment, the method includes administering a pyrophosphate analog. In a second embodiment, the method includes administering a protective agent. Protecting a receptor
5 includes protecting the physical integrity of a receptor and/or the function of a receptor, such as maintaining the ability of the receptor to respond to agonists, to respond to antagonists, to transmit a message to the interior of a cell, or to send a signal within a cell, cell nucleus, or mitochondria.

An embodiment of the invention provides a method for protecting a receptor from
10 free radical damage. Free radicals and other reactive oxygen species (e.g., H_2O_2 , HOCl, and radicals such as O_2^- , sulfur cation, nitric oxide radical, ferryl, peroxy, peroxyxynitrite, thiyl, thiylperoxy, and alkoxy) are highly reactive, and many free radical reactions are highly damaging to cellular components. Free radical reactions can crosslink proteins, mutagenize DNA, and peroxidize lipids. Such reactions can have deleterious effects on
15 cellular receptors. Preferably, the method of the invention includes protection of a receptor, such as a mAChR, or of DNA, RNA, lipids, and proteins necessary for receptor function from deleterious effects.

In another embodiment, the invention provides a method for reducing or eliminating deleterious effects of an endogenous inhibitor found in elevated levels in the
20 brains of Alzheimer's disease patients. This endogenous, low molecular weight inhibitor, as it is known, inhibits agonist and antagonist binding to mAChRs. This inhibitor has a molecular weight of less than 3500 Da and is believed to generate free radicals, in the presence of glutathione or other sulphydryl compounds, that irreversibly inhibit or inactivate the mAChR. The inhibitor also contains free heme, which can
25 generate free radicals, including superoxide radicals, peroxy radicals, and thiyl radicals,

and can cause neurotoxicity. Heme has been shown to damage protein and lipid components of membranes by Vincent (Oxidative Effects of Heme and Porphyrins on Proteins and Lipids, Seminars in Hematology 26(2): 105-113, 1989). Membrane lipid defects have been demonstrated in Alzheimer's disease by Ginsberg et al. (Evidence for a
5 Membrane Lipid Defect in Alzheimer's Disease, Mol. and Chem. Neuropathol. 19: 37-46, 1993). In addition, heme has been proposed to contribute to atherosclerosis by Jacob (Newly recognized causes of atherosclerosis: The role of microorganisms and of vascular iron overload, J. Lab. Clin. Med. 123: 808-816, 1994).

In one embodiment, the method of the invention includes increasing the efficacy
10 of an agent that directly or indirectly affects a mAChR. By way of example, administration of a pyrophosphate analog can increase the efficacy of a muscarinic agonist in the presence of the inhibitor. In another embodiment, the method of the invention includes reducing or eliminating deleterious effects the low molecular weight inhibitor or heme by decreasing or preventing the generation of free radicals or trapping
15 radicals once formed.

Receptors

The invention provides a method for protecting a receptor and/or increasing the efficacy of agents that directly or indirectly affect a receptor. Such receptors include
20 G-protein-coupled receptors, ion-channel-linked receptors and enzyme-linked receptors. Examples include receptors for neurotrophins; neurotransmitters; hormones; steroids; local mediators such as nitric oxide, histamine, and growth factors like PDGF (platelet derived growth factor) and EGF (epithelial growth factor); nucleotides and cyclic nucleotides; calcium and other divalent cations; odorants; antisense oligonucleotides; and
25 the like. Preferably the receptor is a muscarinic receptor. Examples of G-protein

coupled receptors include receptors that respond to odorants, opiates, cannabinoids, glutamate, melatonin, angiotensin II, secretin, vasoactive intestinal peptide (VIP), cholecystokinin (CCK), adrenaline (adrenergic receptors), acetylcholine (muscarinic receptors), ACTH, vasopressin, thrombin, and the like.

5 In one embodiment the invention provides a method for protecting a receptor and/or increasing the efficacy of agents that directly or indirectly affect a receptor in the CNS. In another embodiment, the invention provides a method for protecting a receptor and/or increasing the efficacy of agents that directly or indirectly affect a receptor not in the CNS. Agents whose efficacy are increased by the method of the invention include
10 receptor agonists, allosteric modifiers of receptors, and receptor antagonists.

An embodiment of the invention provides a method for treating or preventing a disease or disorder that relates to or is caused at least in part by dysfunction, alteration, or loss of one or more G-protein coupled receptors. These diseases and disorders include
15 Alzheimer's disease; Parkinson's disease; stroke; multiple sclerosis; ALS; drug addiction, such as opiate addiction or cannabinoid abuse; pain; Sjogren's or dry eye syndrome; heart diseases including congestive heart failure, myocardial infarction, cardiac arrhythmia; cancer; diseases of smooth muscle organs or glands such as irritable bowel syndrome, colitis, hypertension, erectile dysfunction, diabetes, obesity, blood
20 coagulation disorders; and the like.

Muscarinic Receptors

The invention provides a method for protecting a mAChR and/or increasing the efficacy of agents that directly or indirectly affect a mAChR. There are at least five pharmacological classes of mAChRs, including the M1, M2, and M3 muscarinic
25 receptors, and several genetic subclasses including m1, m2, m3, m4, and m5. These

muscarinic receptors are G-protein coupled receptors. Each receptor subtype has its own unique pattern of expression throughout various tissues. As such, dysfunction of each receptor subclass, or combinations thereof, may have deleterious effects leading to a variety of diseases or disorders. The method of the invention can provide protection to a muscarinic receptor in any or several mAChR subclasses, and therefore, can be of benefit to those at risk or suffering from diseases associated with dysfunction of one or more muscarinic receptor subtype. Preferably, the method of the invention provides protection to M1 and M2 muscarinic receptors.

Muscarinic receptors mediate numerous of the inhibitory and excitatory effects of the neurotransmitter acetylcholine in the heart, smooth muscle, blood vessels, glands and in neurons (both presynaptic and postsynaptic) in the autonomic and the central nervous system. Dysfunction of mAChRs thus can contribute to a variety of diseases and/or disorders. Through protection of a mAChR and/or through increasing the efficacy of agents that directly or indirectly affect a mAChR, the method of the invention can provide benefit to subjects suffering from or at risk of a disease or disorder associated with mAChR dysfunction.

An embodiment of the invention provides a method for protecting a mAChR and/or increasing the efficacy of agents that directly or indirectly affect a mAChR in the nervous system of a subject, and therefore, can be of benefit to subjects suffering from or at risk of central nervous system or peripheral nervous system disorders. For example, mAChRs and other receptors are involved in the regulation of the function of cells throughout the CNS. Accordingly, the method of the invention can provide benefit to subjects suffering from or at risk of CNS disorders such as Alzheimer's disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, affective disorders, anxiety

disorders, and/or schizophrenia; nerve damage from cerebrovascular disorders such as stroke, from CNS infections including meningitis and HIV, from tumors of the brain and spinal cord, prion diseases, and CNS disorders resulting from ordinary aging, brain injury, or spinal cord injury. mAChRs are also involved in the regulation of the function of cells in the peripheral nervous system, including the autonomic nervous system. Accordingly, the method of the invention can provide benefit to subjects suffering from or at risk of peripheral nervous system disorders, such as peripheral neuropathy, including that associated with diabetes. For example, in diabetic patients nerves can deteriorate as blood vessels that contain muscarinic or other receptors are lost.

10 Preferably, the method of the invention can benefit a subject suffering from or at risk of Alzheimer's disease.

Another embodiment of the invention provides a method for protecting a mAChR and/or increasing the efficacy of agents that directly or indirectly affect a mAChR not within the nervous system of a subject, and therefore, can be of benefit to subjects suffering from or at risk of disease or disorder outside the nervous system. For example, mAChRs are involved in the regulation (e.g., stimulation or inhibition) of smooth muscle contraction, the regulation of heart rate and cardiac contractility, the regulation of secretion of enzymes or hormones, including the release of amylase from the parotid gland and the release of digestive enzymes and insulin from the pancreas, the regulation of bone growth, and the regulation of iron metabolism. Accordingly, the method of the invention can provide benefit to subjects suffering from or at risk of smooth muscle related disorders such as irritable bowel syndrome, diverticular disease, urinary incontinence, esophageal achalasia, diseases or disorders of the blood vessels (e.g. hypertension), or chronic obstructive airways disease; heart muscle related disorders such as pathologic bradycardia or tachycardia, arrhythmia, flutter or fibrillation; blood related

15
20
25

disorders such as hereditary hematochromatosis; bone disorders such as Paget's disease; cancer; and gland related disorders such as xerostomia, diabetes mellitus, or Sjogren's syndrome or dry eye syndrome which involves decreased secretion of tears by, for example, the lacrimal glands. For example, tear secretion is known to require muscarinic cholinergic stimulation and intact muscarinic receptors. Accordingly, protection of a mACHR and/or increasing the efficacy of an agent that directly or indirectly affects a mACHR can be of benefit to a subject suffering from Sjogren's syndrome or dry eye syndrome.

Protecting a muscarinic receptor can result in enhancing the effectiveness of agents that directly or indirectly affect a mACHR. Useful agents that affect a mACHR include, but are not limited to, anticholinesterase agents, muscarinic agonists, muscarinic antagonists, and other agents useful for treatment of diseases associated with dysfunction of muscarinic receptors, including neurodegenerative and other CNS disorders.

The method of the invention also provides enhanced efficacy of agents that do not act directly, or indirectly, with a mACHR. Such enhanced efficacy can be achieved, for example, through protection of a receptor, preferably a mACHR. Protecting a muscarinic receptor can result in enhancing the efficacy of agents that do not exert their action directly or indirectly on the muscarinic receptor. Such enhanced efficacy can be achieved through desirable effects on cells to which protection of muscarinic receptors provides benefit. Typically, cells that derive benefit from protection of a muscarinic receptor are cells that contain muscarinic receptors. Examples of cells that contain a mACHR include particular neurons, smooth muscle cells, and gland cells.

Cells that lack a muscarinic receptor but which interact with enzymes, hormones, and/or other compounds released from cells with a muscarinic receptor can derive benefit from protecting a muscarinic receptor. Examples of cells lacking a mACHR that can

derive benefit from protecting a mAChR include cells that are presynaptic or postsynaptic relative to cells that contain a mAChR and cells that can interact with enzymes, hormones, and/or other compounds released from cells that contain a mAChR.

Examples of this phenomenon include: Stimulation of m_2 receptors on

5 presynaptic membranes increases the release of acetylcholine which can then stimulate nicotinic receptors on other post synaptic cells. Stimulation of mAChR releases arachidonic acid, which can then affect a variety of other nearby brain cells. Arachidonic acid also increases secretion of amyloid precursor protein. Emmerling, M.R. et al. (1996) Ann. N.Y. Acad. Sci. 777:310-315. Activation of m_1 and m_3 mAChR

10 attenuates release of amyloid B protein. Hung, A.Y. et al. (1993) J. Biol. Chem. 268:22959-22962. Stimulation of mAChR is required for memory and learning which also involve the proper function of noncholinergic cells. Stimulation of mAChR can increase the nitric oxide - cyclic GMP signaling system in neurons (Bauer, M.B. (1994) Neuroscience 62:351-359) and nitric oxide can travel from one cell to another to produce

15 its effects. Stimulation of mAChRs markedly increase hippocampal BDNF and NGF in RNA levels. Once produced, these neurotrophins can produce important effects on other nerve cells in the brain. M. da Penha Berzaghi (1993) J. Neuroscience 13(9) 3818-3826.


By way of further example, some cells containing a mAChR in the pancreas can release insulin. Released insulin can then interact with cells in close proximity to, or at

20 relatively great distances from, the cell from which it was released. Protection of a mAChR on a cell in the pancreas that releases insulin can have beneficial effects on a cell that interacts with insulin. Thus, cells lacking a mAChR can benefit from protection of a mAChR. Similarly, the efficacy of agents acting on cells lacking a mAChR can be enhanced by the method of the invention.

By way of yet further example, stimulation of mAChR in certain brain cells increases potassium ion evoked release of the neurotransmitter dopamine which then goes on to affect other brain cells having dopamine receptors. Joseph, J.A. et al. (1995) Brain Res. 673:195-193. Thus, CNS cells lacking a mAChR can benefit from protection of a mAChR. Similarly, the efficacy of agents acting on CNS cells lacking a mAChR can be enhanced by the method of the invention.

Pyrophosphate Analogs

In one embodiment, the method of the invention provides protection to a receptor and/or increases the efficacy of agents by administering to a subject a pyrophosphate analog. Useful pyrophosphate analogs include compounds of Formula I:




where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²), and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900. Compounds of Formula I in which R¹ is a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²); or R² is H, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl can be referred to as substituted pyrophosphate analogs. Compounds of Formula I can also include substituted pyrophosphate analogs such as dinucleoside-5-5'-pyrophosphates, cyclophosphates of purine and pyrimidine acyclonucleosides. The

compound of Formula I can be any pharmaceutically acceptable salt or basic addition salt. Preferably, X is O, CH₂, NH, or S; R¹ is H; and n is 2-6. More preferably the pyrophosphate analog is pyrophosphate or imidodiphosphate.

Additional preferred compounds of Formula I include pyrophosphate, glycerol pyrophosphate, arachidonylpyrophosphate, imidodiphosphate, serine phosphate, serine imidophosphate, threonine phosphate, threonine imidophosphate, guanylimidodiphosphate and adenylylimidodiphosphate. More preferably compounds of Formula I include pyrophosphate, imidodiphosphate, guanylimidodiphosphate and adenylylimidodiphosphate.

Additional pyrophosphate analogs include compounds of formula II:



where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=3); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group (such as CH₃), or (CH₂)_mNH₂ where m=1-6. Further included are bisphosphonic acids, which are also known as bisphosphonates, where X is preferably RCR¹ and R and R¹ groups are chosen independently from OH, H₂N(CH₂)₂, or CH₃. For example, RCR¹ can be H₂N(CH₂)₂C(OH) or CH₃COH. More specifically, the bisphosphonates include etidronic acid ((1-Hydroxyethylidene)bisphosphonic acid) and pamidronic acid ((3-Amino-1-hydroxypropylidene)bisphosphonic acid) where preferably n=2.

Yet more additional pyrophosphate analogs include substituted pyrophosphate analogs such as inositol diphosphate, inositol triphosphate, inositol tetraphosphate, inositol pentaphosphate, and inositol hexaphosphate.

Such pyrophosphate and imidodiphosphate compounds and the like can be prepared as basic addition salts, such as sodium, potassium, or magnesium salts. It is believed that the use of a basic addition salt, such as a magnesium salt, will reduce the charge and allow for freer movement of the compound throughout the body.

- 5 Pyrophosphate compounds, imidopyrophosphate compounds, and the like can be covalently bound to other phosphates creating polyphosphates or polyimidophosphates. One or more pyrophosphate analogs can be administered in combination. In another embodiment, the pyrophosphate analog can be administered with a protective agent. In another embodiment, the pyrophosphate analog can be administered with a neurologic agent, and optionally with a protective agent.
- 10

Protective Agents

In another embodiment, the invention provides a method for protecting receptors and/or increasing the efficacy of agents by administering to a subject a protective agent.

- 15 Protective agents useful in an embodiment of the method of the invention include a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavonoid; a heme binding compound, such as hemopexin, lipopexin, a lipoprotein, or ApoE-2; and a heme oxygenase, such as heme oxygenase-1 or heme oxygenase-2, or biliverdin reductase, a catalase, a peroxidase, a DNA or RNA vector encoding a biliverdin reductase, a DNA or RNA vector encoding a heme oxygenase (e.g. a DNA or RNA vector encoding a heme oxygenase-1 or a DNA or RNA vector encoding a heme oxygenase-2), a DNA or RNA vector encoding a catalase, a DNA or RNA vector encoding a peroxidase, or a combination thereof Biliverdin reductase is preferably administered with bilirubin, because biliverdin reductase can regenerate bilirubin from biliverdin after bilirubin has
- 20
- 25 been oxidized while functioning as a protective agent. Biliverdin reductase is also

preferably administered in combination with a heme oxygenase. Heme oxygenases include recombinant heme oxygenase. Preferably, a heme oxygenase is a human heme oxygenase.

One or more protective agents can be administered in combination. In another embodiment, one or more protective agents can be administered in combination with one or more pyrophosphate analogs. In another embodiment, one or more protective agents can be administered in with one or more neurologic agents, and optionally with one or more pyrophosphate analogs.

10 **Agents that Directly or Indirectly Affect a mAChR**

The invention also provides a method for enhancing the efficacy of one or more agents that directly or indirectly affect a mAChR. Agents that directly or indirectly affect a mAChR include agents that (1) bind to or interact with a mAChR to either elicit a response transduced through a mAChR or reduce or prevent an agent from binding to or interacting with a mAChR and/or eliciting a signal transduced through a mAChR, (2) alter the concentration of agents that bind to or interact with a mAChR to either elicit a response transduced through a mAChR or reduce or prevent an agent from binding to or interacting with a mAChR and/or eliciting a signal transduced through a mAChR, or (3) modify the ability of agents that bind to or interact with a mAChR to either elicit a response transduced through a mAChR or reduce or prevent an agent from binding to or interacting with a mAChR and/or eliciting a signal transduced through a mAChR. Such agents include anticholinesterase agents, muscarinic agonists, muscarinic antagonists, and allosteric modifiers of muscarinic receptors. Preferably, the invention provides a method for enhancing the efficacy of agents that either directly or indirectly elicit a

response through a mAChR. Most preferably, the method of the invention enhances the efficacy of a muscarinic agonist or an anticholinesterase agent.

Muscarinic receptor agonists directly elicit a response through a mAChR by binding to and transducing a signal through a mAChR. Preferred muscarinic agonists include acetylcholine, Xanomeline, and the like.

A muscarinic antagonist is an agent that is capable of partially or completely inhibiting, or reversing, the effect of a muscarinic agonist on a mAChR. Examples of muscarinic antagonists include atropine, N-methyl-scopolamine, quinuclidinyl benzilate, pirenzepine, and the like.

Cholinesterase hydrolyzes the neurotransmitter acetylcholine and provides one of the mechanisms responsible for rapid depletion of acetylcholine from the synaptic cleft. Anticholinesterase agents inhibit cholinesterase activity and as a result increase the concentration of acetylcholine in the synaptic cleft and prolong the duration of which acetylcholine remains in the synaptic cleft. Anticholinesterase agents can thus indirectly affect a mAChR by increasing concentrations of and prolonging the effective duration of acetylcholine in the synaptic cleft. Anticholinesterase agents can also interact directly with cholinergic receptors, including mAChRs; with sodium and potassium ion channels; and effect the uptake, synthesis and release of neurotransmitters. Preferred anticholinesterase agents include Aricept, Exelon, Metrifonate, and the like.

An allosteric modifier of a mAChR binds or interacts with a site other than the agonist binding site of a mAChR and modifies the ability of an agonist or an antagonist to elicit or inhibit, respectively, a response transduced through a muscarinic receptor, without itself eliciting a response. Suitable allosteric modifiers of a muscarinic agonist include gallamine and dynorphin. Preferred allosteric modifiers of a mAChR include gallamine and dynorphin.

Neurologic Agents

The invention also provides a method for increasing the efficacy of a neurologic agent in a subject in need thereof. In a first embodiment, this method includes
5 administering a pyrophosphate analog. In this embodiment, the increased efficacy of the neurologic agent preferably results from protection of a muscarinic receptor caused or induced by the pyrophosphate analog. In a second embodiment, this method includes administering a protective agent. In this second embodiment, the increased efficacy of the neurologic agent preferably results from protection of a muscarinic receptor caused or
10 induced by the protective agent. In each embodiment, the subject preferably is concurrently receiving, has recently received, or will soon receive a neurologic agent.

A neurologic agent promotes nerve cell growth and survival or augments the activity of functioning cells. Among those agents that are preferred are cholinergic agonists, allosteric modifiers of a mAChR, cholinesterase inhibitors, or neurotrophic and
15 neuritogenic factors that are similar to naturally occurring nerve growth promoting substances. Among the preferred neurologic agents are gangliosides (such as GM-I ganglioside), phosphatidylserine (PS), nerve growth factor (NGF), neurotrophins 3, 4, and/or 5 (NT-3, NT-4 and/or NT-5) brain-derived neurotrophic factor (BDNF), fibroblast growth factors (FGFs, e.g., basic fibroblast growth factor), insulin, insulin-like growth
20 factors (IGF-1 and/or IGF-2), ciliary neurotrophic factor (CNTF), transforming growth factors, epidermal growth factors, activity-dependent growth factor, platelet derived growth factor, neurokinin, glia-derived neurotrophic factor (GDNF), glia-derived nexin, and cholinergic enhancing factors such as phosphoethanolamine and thyroid hormone T₃, and DNA or RNA vectors or plasmids that encode one or more protein neurologic

agents or nerve growth promoting factors. Plasmids and vectors for delivery of a coding sequence to a mammalian tissue are known to those of skill in the art.

Metal Diseases

5 The method of the invention can treat or prevent diseases or disorders caused or induced by metals, such as cancer and poisoning. Such metals can include As, Co, Cr, Ni, Hg, Pb, Fe, Cu, V, and Cd. That is, the method of the invention can treat or prevent poisoning by (for example) lead or mercury and also excessive iron toxicity. In one embodiment, the method of the invention can treat or prevent CNS diseases or disorders
10 caused or induced by metals. In another embodiment, the method of the invention can treat or prevent diseases or disorders not of the CNS but caused or induced by metals, such as heart disease, blood vessel disease, and gland disease. In another embodiment, the method of the invention reduces poisoning of a subject by at least one metal. In another embodiment, the method of the invention protects a subject from at least one
15 carcinogenic metal. In another embodiment, the method of the invention reduces toxic actions of metal ions in a subject, particularly toxic actions due to Fe^{++} , Hg^{++} , Cd^{++} , Cu^{++} , As^{+++} , and Pb^{++} ions.

Administering Agents

20 Administering compounds according to the method of the invention can include formulating the compounds or compositions as pharmaceutical compositions and administering the pharmaceutical compositions to a mammalian host, including a human patient, in a variety of forms adapted to the chosen route of administration. The compounds are preferably administered in combination with a pharmaceutically
25 acceptable carrier. The compounds can be administered at one of a variety of doses

sufficient to provide an effective amount at the desired point of action of the agent.

Doses for humans and other mammals can range from about 0.001 mg/kg to about 100 mg/kg, preferably from about 0.01 mg/kg to about 10 mg/kg, preferably from about 0.1 mg/kg to about 1-10 mg/kg.

5 A related use of the methods of the invention is to protect pharmacological agents in formulation. The pharmacological agents may be for therapeutic, diagnostic, or other purposes.

 The compounds can be administered by known techniques, such as orally, intranasally, parentally (including subcutaneous injection, intravenous, intramuscular, 10 intrasternal or infusion techniques), by inhalation spray, dermally, transdermally, intrathecal, intracerebroventricular, buccal, sublingual, topically, by absorption through a mucous membrane or through the skin, or rectally, in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants or vehicles. Pharmaceutical compositions of the invention can be in the form of 15 suspensions or tablets suitable for oral administration, nasal sprays, eye drops, nose drops, creams, sterile injectable preparations, such as sterile injectable aqueous or oleagenous suspensions or suppositories.

 Controlled or sustained release systems can also be employed. For example, compositions can include a polymer or other substance that enhances controlled or 20 sustained release. Controlled or sustained release systems can include a polymer disk, such as evac disks, microspheres, and copolymers. Preferred controlled release polymers are poly(lactide:glycolide) and poly(ethylene-co-vinyl acetate).

 For oral administration as a suspension, the compositions can be prepared according to techniques well known in the art of pharmaceutical formulation. The 25 compositions can contain microcrystalline cellulose for imparting bulk, alginic acid or

WO 01/23021
sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents. As immediate release tablets, the compositions can contain microcrystalline cellulose, starch, magnesium stearate and lactose or other excipients, binders, extenders, disintegrants, diluents and lubricants known in the art.

5 In addition to the typical pharmacological methods for oral administration, the agents employed in the methods of the invention can be administered as a component of a nutritional or food supplement. The nutritional or food supplement can also include any other ingredients typical of a nutritional or food supplement, such as flavorings, stabilizers, and the like.

10 For administration by inhalation or aerosol, the compositions can be prepared according to techniques well known in the art of pharmaceutical formulation. The compositions can be prepared as solutions in saline, using benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons or other solubilizing or dispersing agents known in the art.

15 For administration as injectable solutions or suspensions, the compositions can be formulated according to techniques well-known in the art, using suitable dispersing or wetting and suspending agents, such as sterile oils, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

 For intranasal administration, the compositions can be formulated according to
20 techniques well known in the art. The means of applying a pharmaceutical composition intranasally can be in a variety of forms such as a powder, spray or nose drops.

 For transdermal administration, the compositions can be formulated according to techniques well known in the art. Delivery of the composition through the skin can be accomplished by delivery means well known in the art, including transdermal patch, an
25 ointment, an iontophoretic patch or device, and the like.

For rectal administration as suppositories, the compositions can be prepared by mixing with a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ambient temperatures, but liquefy or dissolve in the rectal cavity to release the drug.

- 5 Preferred administration routes include orally, parenterally, as well as intravenous, intramuscular or subcutaneous routes.

Intraocular administration through the use of an ointment or eye drops is preferred for treatment of a glandular disease or disorder of the eye, such as a lachrymal gland disease disorder like Sjogren's Syndrome or dry eye syndrome.

- 10 Solutions or suspensions of the compounds can be prepared in water, isotonic saline (PBS) and optionally mixed with a nontoxic surfactant. Dispersions may also be prepared in glycerol, liquid polyethylene, glycols, DNA, vegetable oils, triacetin and mixtures thereof Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

- 15 The pharmaceutical dosage form suitable for injection or infusion use can include sterile, aqueous solutions or dispersions or sterile powders including an active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle
- 20 can be a solvent or liquid dispersion medium including, for example, water, ethanol, a polyol such as glycerol, propylene glycol, or liquid polyethylene glycols and the like, vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size, in the case of dispersion, or by the use of
- 25 nontoxic surfactants. The prevention of the action of microorganisms can be

WO 01/021952
accomplished by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be desirable to include isotonic agents, for example, sugars, buffers, or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the
5 inclusion in the composition of agents delaying absorption--for example, aluminum monostearate hydrogels and gelatin.

Sterile injectable solutions are prepared by incorporating the compounds in the required amount in the appropriate solvent with various other ingredients as enumerated above and, as required, followed by filter sterilization. In the case of sterile powders for
10 the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

15 **Administering Agents to the Brain**

Administering agents, e.g. a protective agent, a pyrophosphate analog, an agent that directly or indirectly affects a mAChR, and/or a neurologic agent, according to the method of the invention includes administering agents to a mammalian host in a manner that allows the agents to exert their effect in the CNS. Many agents useful for the
20 method of the invention can be absorbed into the blood stream and readily cross the blood brain barrier.

However, some agents useful for the method of the invention cannot pass, or have difficulty passing, the blood brain barrier. Such agents can be administered as "prodrugs" which can cross the blood brain barrier, and upon or after entry into the CNS,

the prodrug is converted to the active agent. Agents that can cross the blood brain barrier without difficulty can also be administered as prodrugs.

To deliver the agent to the CNS, the agent alone or in combination with other substances as a pharmaceutical composition may be administered to the spinal cord and
5 to the cerebral vesicles according to intrathecal and intracerebrovascular administration methods known in the art. Such pharmaceutical compositions can also be administered to the nasal cavity, under the tongue, or onto the eye. The composition may be dispensed intranasally, sublingually, or conjunctivally as a powdered or liquid nasal spray, nose drops, a gel or ointment, through a tube or catheter, by syringe, by packtail, by pledget,
10 or by submucosal infusion. The agent may be combined with a polymer or other substance that enhances controlled or sustained release of the agent. In particular, agents can be delivered to the brain by intranasal administration as described in X.-Q. Chen et al. (1998) J. Alzheimer's Disease 1:35-44 and W.H. Frey II et al. (1997) Drug Delivery 4:87-92; the disclosures of which are incorporated herein by reference.

15 The optimal concentration of the active agent will necessarily depend upon the specific agent used, the characteristics of the patient and the nature of the disease or condition for which the treatment is to be used.

The carrier of the composition may be any material which is otherwise pharmaceutically-acceptable and compatible with the active ingredients of the
20 composition. Where the carrier is a liquid, it is preferred that the carrier is hypotonic or isotonic with nasal, oral, or conjunctival fluids and have a pH within the range of 4.5-7.5. Where the carrier is in powdered form, it is preferred that the carrier is also within an acceptable non-toxic pH range.

The pharmaceutical composition may be formulated as a powder, granules,
25 solution, ointment, cream, aerosol, powder, drops, or a controlled or sustained release

composition such as a polymer disk. The solution may be sterile, isotonic or hypotonic, and otherwise suitable for administration by injection or other means. In addition to the agent, the solution may contain appropriate adjuvants, buffers, preservatives and salts. The powder or granular forms of the pharmaceutical composition may be combined with
5 a solution and with diluting, dispersing or surface active agents. Solutions such as nose or eye drops may contain an antioxidant, a buffer, and the like. Further controlled release polymers may be used to regulate the delivery of the agent.

The present invention may be better understood with reference to the following examples. These examples are intended to be representative of specific embodiments of
10 the invention, and are not intended as limiting the scope of the invention.

EXAMPLES

Example 1 - Protection of Muscarinic Acetylcholine Receptor (mAChR) in Cell Free

Systems

15 Materials and Methods

Membrane mAChR Preparation

Membranes rich in mAChRs were prepared by a modification of the method used by Marks and Collins (Characterization of nicotine binding in mouse brain and comparison with the binding of α -bungarotoxin and quinuclidinyl benzilate, Mol.
20 Pharmacol. 22:544-564, 1982). Gray matter from nondemented adult human frontal cortex was homogenized in 9 vol of 50 mM Tris-HCl, pH 7.4, using 5 passes of a glass/Teflon motor-driven homogenizer. The homogenate was centrifuged at 27 000 x g for 20 mm at 4°C, and the subsequent pellet resuspended in 9 vol of cold deionized water with 5 passes of the homogenizer. The resuspension was incubated at 37°C for 5 min,
25 then was centrifuged as before. The resulting pellet was resuspended, incubated and

centrifuged again as above. The final pellet was weighed, resuspended at 15% w/v in 50 mM Tris-HCl buffer, aliquoted in small portions, flash-frozen in liquid nitrogen and stored at -70°C for subsequent assays to determine content and binding capacity. Before use in binding assays, the thawed membrane preparation was briefly rehomogenized with 10 passes in a glass/glass homogenizer. A typical mAChR membrane preparation bound 300 pmol [³H] quinuclidinyl benzilate ([³H]QNB)/g protein.

Inhibitor Preparation

Gray matter obtained from the frontal cortex of cases with AD was homogenized in 9 vol of 1% trifluoroacetic acid (TFA) for 40 s at 4°C in a Waring blender, then centrifuged at 1200 x g for 10 min at 4°C. The resulting supernatant fraction was centrifuged at 11 000 x g for 100 min at 4°C. The 11 000 x g supernatant fraction was centrifuged at 100 000 x g for 100 min at 4°C, then the 100 000 x g supernatant fraction was concentrated using a SpeedVac and resuspended in 0.1% TFA to half the original tissue volume. The 100 000 x g supernatant fraction was transferred to a Spectra/Por 3 dialysis membrane bag (3500 dalton cutoff), and dialyzed against 20 vol of 0.1% TFA at 4°C for 24 h with gentle stirring. The resulting <3500 Da fraction (dialysate) was concentrated by SpeedVac to half the original tissue volume. The <3500 Da fraction (endogenous inhibitor) was then frozen in liquid nitrogen and stored at -70°C for subsequent assays to determine protein count and inhibitor activity. Protein activity was measured using the bicinchononic acid (BCA) protein assay method, essentially as described by Smith et al. (Measurement of protein binding using bicinchonic acid, Ann. Biochem. 150: 76-85, 1985). A typical inhibitor preparation contained about 4 mg/ml protein and approximately twice the concentration of inhibitor found in the original tissue.

Inhibitor Activity Assay

Inhibitor activity was measured using a modification of the method of Fields et al. (Cardiac muscarinic receptors, J. Biol. Chem. 253:3251-3258, 1978) to assess the binding of [3 H]QNB, a mAChR antagonist, or [3 H]-oxotremorine M, a mAChR agonist. In general, binding conditions consisted of 50 mM Tris-HCl, pH 7.4 at 37°C, 10 mM reduced glutathione (GSH), 75 μ /ml membrane and 2×10^{-10} M [3 H]QNB or 3nM [3 H]-oxotremorine M, with and without addition of inhibitor. To control for non-specific binding, 12.5 μ M atropine sulfate (a mAChR antagonist) was added to several tubes. Subtracting nonspecific binding from total binding yielded specific binding.

Pyrophosphate, imidodiphosphate, adenylylimidodiphosphate, guanylimidodiphosphate, and tripolyphosphate were dissolved in distilled water. Bilirubin, biliverdin, and heme were dissolved in DMSO. Carnosol, myricetin, and quercetin were dissolved in ethanol. Catalase and peroxidase were dissolved in an aqueous, preferably buffered, solution.

Enough water was added to all other reaction components in each tube to make 4 ml total. The binding reaction was initiated by adding [3 H]QNB or [3 H]-oxotremorine-M, mixing the tubes briefly, and then incubating the tubes at 37°C for [3 H]QNB or at room temperature for [3 H]-oxotremorine-M. The reaction time for [3 H]QNB was one hour in most experiments. In some experiments, the mAChR was preincubated with either the endogenous LMW inhibitor or heme plus peroxide in the presence or absence of the therapeutic agent being tested. The effect of the therapeutic agent on receptor function was then assessed in a binding assay, which for [3 H]QNB was conducted at 37°C for 40 min, and for [3 H]-oxotremorine-M was conducted at room temperature for 20 min. After 60 min, the binding reaction was terminated by adding 5 ml of cold 50

mM Tris buffer, pH 7.4, to each tube and chilling the tubes in an ice bath. The tube contents and one 15 ml rinse of cold 50 mM Tris buffer, pH 7.4, were filtered through Whatman GF/B glass fiber filters using a Brandel harvester. The filters were placed in Optiflow scintillation flour and counted in a Beckman LS-6500 scintillation counter set for tritium detection.

Results

The data resulting from the methods presented above and the results presented in Figures 1-20 are discussed in more detail below.

Pyrophosphate:

Pyrophosphate protects the mAChR from inactivation by the LMW inhibitor or by the combination of heme and peroxide. Pyrophosphate protected the receptor from both loss of antagonist (^3H -QNB) binding (Figures 1 and 2) and agonist (^3H -Oxotremorine M) binding (Figure 3). Approximately 1 μM pyrophosphate provides 50% protection. Pyrophosphate also protects the mAChR from damage by PbCl_2 . Approximately 57 μM pyrophosphate provides 50% protection (Figure 20).

Imidodiphosphates:

Imidodiphosphate (Figure 4), guanylimidodiphosphate (Figure 5), and adenylylimidodiphosphate (Figure 6) all protect the mAChR from inactivation by the LMW inhibitor.

Polyphosphates:

WO 01/82932
Polyphosphates, such as tripolyphosphate (Figure 7), protect the mAChR from inactivation by the LMW inhibitor.

Bisphosphonates:

5 Bisphosphonates, such as pamidronate (Figure 19), protect the mAChR from inactivation by the LMW inhibitor.

Bilirubin and Biliverdin:

10 Bilirubin protects the mAChR from inactivation by the LMW inhibitor or by the combination of heme and peroxide. Approximately 0.7 μ M bilirubin provides 50% protection of the receptor from loss of antagonist (3 H-QNB) binding (Figures 8 and 9) and 1.9 μ M provides 50% protection from loss of agonist (3 H-Oxotremorine M) binding (Figure 10).

biliverdin at 3 μ M provides 50% protection of the mAChR (Figure 11).

15

Carnosol, Quercetin, and Myricetin:

Carnosol (Figure 12), quercetin (Figure 13), and myricetin (Figure 14) all protected the mAChR from inactivation. Carnosol provided 100% protection at 1 μ M. while quercetin and myricetin provided 50% protection at 0.24 μ M and 0.4 μ M
20 respectively.

Catalase and Peroxidase

Catalase protected the mAChR from inactivation by the LMW inhibitor or by the combination of heme and peroxide (Figures 15 and 16, respectively). As little as 0.34
25 units/mL of catalase provided 50% protection from inactivation by heme and peroxide.

A peroxidase, specifically glutathione peroxidase, protected the mAChR from inactivation by the LMW inhibitor or by the combination of heme and peroxide (Figures 17 and 18, respectively). Glutathione peroxidase at 0.5 units/mL provided 71% protection from inactivation by heme and peroxide.

5

Conclusion

The results indicate that pyrophosphate, imidodiphosphates, polyphosphates, bisphosphonates, bilirubin, biliverdin, carnosol, quercetin and myricetin protect a receptor and increase the ability of agents to bind a receptor. Particularly, the results demonstrate the ability of these agents to protect a muscarinic receptor from the effects of endogenous LMW inhibitor, heme and metals and increase the ability of muscarinic agonists and antagonists to bind a mAChR, suggesting that these agents can be used effectively to protect other receptors and increase the efficacy of other agents.

Because the mAChR is essential for memory and learning, the specific demonstration that these agents can protect the human brain mAChR from inactivation and increase agonist binding indicates that these agents have therapeutic potential for the treatment of cognitive and memory disorders including those associated with aging, such as Alzheimer's disease.

20

Example 2- Protection of the mAChR in Cell Culture

Various systems for determining the protection of a mAChR in cell culture are known in the art. Such cell culture systems can be used to determine if mAChR is protected from a damaging agent or condition according to the method of the invention. For example, by administering a mAChR antagonist or agonist alone or in combination with one or more protective agent and/or one or more pyrophosphate analog, one of skill

25

WO 01/02932 PC1/US01/13931
in the art can determine if a mAChR is protected by the one or more protective agent
and/or one or more pyrophosphate analog.

Example 3- Protection of the mAChR Receptor in Animals

5 Various systems for determining the protection of a mAChR in animals are
known in the art. Such animal systems can be used to determine if mAChR is protected
according to the method of the invention. For example, by administering a mAChR
antagonist or agonist alone or in combination with one or more protective agents and/or
one or more pyrophosphate analogs, one of skill in the art can determine if a mAChR is
10 protected by the one or more protective agents and/or one or more pyrophosphate
analog.

Example 4- Increased Efficacy of Neurologic Agents in Model Systems

 Various models systems for determining the efficacy of neurologic agents are
15 known in the art. Such model systems can be used to determine if the efficacy of a
neurologic agent is increased by the method of the invention. For example, by
administering one or more neurologic agents alone or in combination with one or more
protective agents and/or one or more pyrophosphate analogs, one of skill in the art can
determine if the efficacy of the one or more neurologic agents is increased when
20 administered with one or more protective agents and/or one or more pyrophosphate
analog within the parameters of the model system according to techniques known in the
art.


 It should be noted that, as used in this specification and the appended claims, the
singular forms “a,” “an,” and “the” include plural referents unless the content clearly
25 dictates otherwise. Thus, for example, reference to a composition containing “a

compound” includes a mixture of two or more compounds. It should also be noted that the term “or” is generally employed in its sense including “and/or” unless the content clearly dictates otherwise.

5 All publications and patent applications in this specification are indicative of the level of ordinary skill in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated by reference.

10 The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many variations and modifications may be made while remaining within the spirit and scope of the invention.


1. A method of protecting a tissue component in a subject in need thereof, comprising administering to the subject at least one pyrophosphate analog comprising the structure:



where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;

10

or the structure:



where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;


or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate;

15

or a pharmaceutically acceptable salt thereof.


2. The method of claim 1, wherein the tissue component is at least one of a receptor, a protein, a lipid, a nucleic acid, a carbohydrate, a hormone, a vitamin, and a cofactor.
- 20

3. The method of claim 2, in which the tissue component is a receptor for a neurotransmitter, a neuropeptide, a neurotrophin, a growth factor, a steroid, a histamine, a purine, a benzodiazepine, arachidonic acid, nitric oxide, carbon monoxide, an odorant, or an ion channel.
- 5 4. A method of protecting tissue from oxidative stress, comprising administering at least one pyrophosphate analog comprising the structure:



where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;

or the structure:




where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;

or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate;


or a pharmaceutically acceptable salt thereof.

- 20 5. A method of increasing the efficacy of an agent that directly or indirectly affects a receptor in a subject in need thereof, comprising administering the agent, and administering at least one pyrophosphate analog comprising the structure:



where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;

or the structure:




where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;

or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate;


or a pharmaceutically acceptable salt thereof.

6. The method of claim 5, in which the receptor is a receptor for a neurotransmitter, a neuropeptide, a neurotrophin, a growth factor, a steroid, a histamine, a purine, a benzodiazepine, arachidonic acid, nitric oxide, carbon monoxide, an odorant, or an ion channel.
7. The method of claim 6, in which the receptor is one of muscarinic acetylcholine, nicotinic acetylcholine, an opiate, a catecholamine, serotonin, glutamate, aspartate, cannabinoid, gamma aminobutyric acid, or glycine.
8. The method of claim 6, in which the agent directly or indirectly affects a mAChR.

9. The method of claim 8, wherein the agent that directly or indirectly affects a mAChR comprises an anticholinesterase agent, a neurologic agent, or a muscarinic receptor agonist.
10. The method of claim 8, in which the agent that directly or indirectly affects a mAChR comprises Xanomeline.
11. The method of claim 8, wherein the agent that directly or indirectly affects a mAChR comprises donepezil, rivastigmine, galanthamine, metrifonate, or a combination thereof.
12. A method of protecting a subject from at least one carcinogenic metal, comprising administering to the subject at least one pyrophosphate analog comprising the structure:



- where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;




or the structure:

- where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;

or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate; or a pharmaceutically acceptable salt thereof.


- 5 13. The method of claim 12, in which the carcinogenic metal is selected from the group consisting of arsenic, cadmium, cobalt, nickel, lead, and chromium.
14. A method of reducing poisoning of a subject by at least one metal, comprising administering to the subject at least one pyrophosphate analog comprising the structure:



where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;

15

or the structure:




where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;

or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate; or a pharmaceutically acceptable salt thereof.


20

15. The method of claim 14, in which the metal is selected from the group consisting of iron, copper, mercury, lead, cadmium, vanadium and their alloys.
16. A method of treating bacterial, fungal, algo, or algae infections in a subject in need thereof, comprising administering to the subject at least one pyrophosphate analog comprising the structure:




- where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;

or the structure:




- where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;
- 15 or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate;
- or a pharmaceutically acceptable salt thereof.
17. The method of claim 16, in which the subject is a plant.
- 20 18. The method of claim 17, in which the pyrophosphate analog comprises imidodiphosphate.
19. The method of claim 16, in which the subject is an animal or mammal.

20. A method of reducing toxic actions of metal ions in a subject in need thereof, comprising administering to the subject at least one pyrophosphate analog comprising the structure:



- 5 where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;


- 10 or the structure:



where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;


- 15 or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate;
- or a pharmaceutically acceptable salt thereof.

21. The method of claim 20, in which the metal ion is selected from the group consisting of Fe⁺⁺, Hg⁺⁺, Cd⁺⁺, Cu⁺⁺, As⁺⁺⁺, and Pb⁺⁺.
22. A method of protecting a pharmacological agent in a formulation, comprising
- 20 combining the agent with at least one pyrophosphate analog comprising the structure:



where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;

or the structure:




where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;

or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate;


or a pharmaceutically acceptable salt thereof.

23. The method of claim 22, in which the pharmacological agent is a therapeutic agent.
24. The method of claim 22, in which the pharmacological agent is a diagnostic agent.
25. A method of increasing efficacy of a neurologic agent in a subject in need thereof, comprising administering to the subject the neurologic agent, and administering at least one pyrophosphate analog comprising the structure:



where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²) and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900;

or the structure:



where n=2-4; X is O; RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, or (CH₂)_mNH₂ where m=1-6;

or a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, a pyrimidine acyclonucleoside, an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate;

or a pharmaceutically acceptable salt thereof.


26. The method of claim 25, wherein the neurologic agent comprises a ganglioside, a phosphatidylserine, a nerve growth factor, a neurotrophin, a brain-derived neurotrophic factor, a fibroblast growth factor, an insulin, an insulin-like growth factor, a transforming growth factor, an epidermal growth factor, a platelet-derived growth factor, a neurokinin, activity-dependent neurotrophic factor, a ciliary neurotrophic factor, a glia-derived neurotrophic factor, a glia-derived nexin, a cholinergic enhancing factor, an antisense oligonucleotide, a DNA or RNA vector or plasmid that encodes one or more protein neurologic agents or nerve growth promoting factors, or a combination thereof.

27. The method of claim 26, wherein the ganglioside comprises GM-1 ganglioside.

28. The method of claim 26, wherein the neurotrophin comprises neurotrophin 3, neurotrophin 4, neurotrophin 5, or a combination thereof.
29. The method of claim 26, wherein the fibroblast growth factor comprises basic fibroblast growth factor, or acidic fibroblast growth factor.
- 5 30. The method of claim 26, wherein the insulin-like growth factor comprises insulin-like growth factor-I, insulin-like growth factor-2, or a combination thereof.
31. The method of claim 26, wherein the cholinergic enhancing factor comprises ethanolamine, thyroid hormone T.3, gallamine or a combination thereof.
32. The method of any of claims 1, 4, 5, 12, 14, 16, 20, 22, or 25, wherein the
10 pyrophosphate analog comprises pyrophosphate, imidodiphosphate, guanylimidodiphosphate, adenylylimidodiphosphate, or a bisphosphonate.
33. The method of claim 32, wherein the pyrophosphate analog comprises etidronic acid, pamidronic acid, or a combination thereof.
34. The method of any of claims 1, 4, 5, 12, 14, 16, 20, 22, or 25, wherein the subject
15 suffers from at least one of cancer; neuropathies, diseases, or disorders of the heart, smooth muscles, blood, blood vessels, glands, or bones; hypertension; myocardial infarction, ischemic heart disease, or congestive heart failure; irritable bowel syndrome; diverticular disease; urinary incontinence; esophageal achalasia; chronic obstructive airways disease; cardiac arrhythmia; xerostomia; diabetes
20 mellitus; Sjogren's syndrome; Paget's disease; hereditary hematochromatosis; or a non-CNS disorder resulting from normal aging.
35. The method of any of claims 1, 4, 5, 12, 14, 16, 20, 22, or 25, wherein the subject suffers from at least one of a neurologic disorder and a psychiatric disorder.
36. The method of claim 35, wherein the subject suffers from at least one of
25 Alzheimer's disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, an affective disorder, an anxiety disorder, schizophrenia, cell damage, nerve damage, a CNS infection, a tumor of the brain, a tumor of the spinal cord, a stroke in the brain, a stroke in the spinal cord, a prion disease, a CNS disorder

resulting from ordinary aging, a brain injury, a spinal cord injury; or a non-CNS disorder resulting from normal aging.

37. The method of any of claims 1, 4, 5, 12, 14, 16, 20, 22, or 25, further comprising combining any pyrophosphate analog with at least one of a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavonoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
38. The method of claim 37, further comprising administering heme oxygenase, a vector encoding a heme oxygenase, heme oxygenase-1, a vector encoding a heme oxygenase-1, heme oxygenase-2, a vector encoding a heme oxygenase-2, a biliverdin reductase, a vector encoding a biliverdin reductase, a catalase, a vector encoding a catalase, a peroxidase, a vector encoding a peroxidase, or a combination thereof.
39. The method of claim 37, further comprising administering a heme binding protein.
40. The method of claim 39, wherein the heme binding protein comprises hemopexin, a lipoprotein, or a combination thereof.
41. The method of any of claims 1, 4, 5, 12, 14, 16, 20, 22, or 25, further comprising administering heme oxygenase, a vector encoding a heme oxygenase, heme oxygenase-1, a vector encoding a heme oxygenase-1, heme oxygenase-2, a vector encoding a heme oxygenase-2, a biliverdin reductase, a vector encoding a biliverdin reductase, a catalase, a vector encoding a catalase, a peroxidase, a vector encoding a peroxidase, or a combination thereof.
42. The method of any of claims 1, 4, 5, 12, 14, 16, 20, 22, or 25, further comprising administering a heme binding protein.
43. The method of claim 42, wherein the heme binding protein comprises hemopexin, a lipoprotein, or a combination thereof.
44. The method of any of claims 1, 4, 5, 12, 14, 16, 20, 22, or 25, in which the compound is:



and n is 1-6.

45. A method of protecting a tissue component in a subject in need thereof, comprising administering to the subject a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavinoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
46. The method of claim 45, wherein the tissue component is at least one of a receptor, a protein, a lipid, a nucleic acid, a carbohydrate, a hormone, a vitamin, and a cofactor.
47. The method of claim 46, in which the tissue component is a receptor for a neurotransmitter, a neuropeptide, a neurotrophin, a growth factor, a steroid, a histamine, a purine, a benzodiazepine, arachidonic acid, nitric oxide, carbon monoxide, an odorant, or an ion channel.
48. A method of protecting tissue from oxidative stress, comprising administering a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavinoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
49. A method of increasing the efficacy of an agent that directly or indirectly affects a receptor in a subject in need thereof, comprising administering the agent, and administering a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavinoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
50. The method of claim 49, in which the receptor is a receptor for a neurotransmitter, a neuropeptide, a neurotrophin, a growth factor, a steroid, a histamine, a purine, a benzodiazepine, arachidonic acid, nitric oxide, carbon monoxide, an odorant, or an ion channel.

51. The method of claim 50, in which the receptor is one of muscarinic acetylcholine, nicotinic acetylcholine, an opiate, a catecholamine, serotonin, glutamate, aspartate, cannabinoid, gamma aminobutyric acid, or glycine.
52. The method of claim 49, in which the agent directly or indirectly affects a mAChR.
53. The method of claim 52, wherein the agent that directly or indirectly affects a mAChR comprises an anticholinesterase agent, a neurologic agent, or a muscarinic receptor agonist.
54. The method of claim 52, in which the agent that directly or indirectly affects a mAChR comprises Xanomeline.
55. The method of claim 52, wherein the agent that directly or indirectly affects a mAChR comprises donepezil, rivastigmine, galanthamine, metrifonate, or a combination thereof.
56. A method of protecting a subject from at least one carcinogenic metal, comprising administering to the subject a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavonoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
57. The method of claim 56, in which the carcinogenic metal is selected from the group consisting of arsenic, cadmium, cobalt, nickel, lead, and chromium.
58. A method of reducing poisoning of a subject by at least one metal, comprising administering to the subject a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavonoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
59. The method of claim 56, in which the metal is selected from the group consisting of iron, copper, mercury, lead, cadmium, and their alloys.
60. A method of treating bacterial, fungal, algo, or algae infections in a subject in need thereof, comprising administering to the subject a bilirubin, biliverdin,

carosol, quercetin, myricetin, a bioflavinoid, a combination thereof, or a pharmaceutically acceptable salt thereof.

61. The method of claim 60, in which the subject is a plant.
62. The method of claim 60, in which the subject is an animal or mammal.
- 5 63. A method of reducing toxic actions of metal ions in a subject in need thereof, comprising administering to the subject a bilirubin, biliverdin, carosol, quercetin, myricetin, a bioflavinoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
- 10 64. The method of claim 63, in which the metal ion is selected from the group consisting of Fe^{++} , Hg^{++} , Cd^{++} , Cu^{++} , As^{+++} , and Pb^{++} .
65. A method of protecting a pharmacological agent in a formulation, comprising combining the agent with at least one of a bilirubin, biliverdin, carosol, quercetin, myricetin, a bioflavinoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
- 15 66. The method of claim 65, in which the pharmacological agent is a therapeutic agent.
67. The method of claim 65, in which the pharmacological agent is a diagnostic agent.
- 20 68. A method of increasing efficacy of a neurologic agent in a subject in need thereof, comprising administering to the subject the neurologic agent, and administering a bilirubin, biliverdin, carosol, quercetin, myricetin, a bioflavinoid, a combination thereof, or a pharmaceutically acceptable salt thereof.
- 25 69. The method of claim 68, wherein the neurologic agent comprises a ganglioside, a phosphatidylserine, a nerve growth factor, a neurotrophin, a brain-derived neurotrophic factor, a fibroblast growth factor, an insulin, an insulin-like growth factor, a transforming growth factor, an epidermal growth factor, a platelet-derived growth factor, a neurokine, activity-dependent neurotrophic factor, a ciliary neurotrophic factor, a glia-derived neurotrophic factor, a glia-derived


nexin, a cholinergic enhancing factor, an antisense oligonucleotide, a DNA or RNA vector or plasmid that encodes one or more protein neurologic agents or nerve growth promoting factors, or a combination thereof.

70. The method of claim 69, wherein the ganglioside comprises GM-1 ganglioside.
- 5 71. The method of claim 69, wherein the neurotrophin comprises neurotrophin 3, neurotrophin 4, neurotrophin 5, or a combination thereof.
72. The method of claim 69, wherein the fibroblast growth factor comprises basic fibroblast growth factor, or acidic fibroblast growth factor.
73. The method of claim 69, wherein the insulin-like growth factor comprises insulin-like growth factor-I, insulin-like growth factor-2, or a combination thereof.
- 10 74. The method of claim 69, wherein the cholinergic enhancing factor comprises ethanolamine, thyroid hormone T.3, gallamine or a combination thereof.
75. The method of any of claims 45, 47, 48, 49, 56, 58, 60, 63, 65, or 68, wherein the subject suffers from at least one of cancer; neuropathies, diseases, or disorders of the heart, smooth muscles, blood, blood vessels, glands, or bones; hypertension; myocardial infarction, ischemic heart disease, or congestive heart failure; irritable bowel syndrome; diverticular disease; urinary incontinence; esophageal achalasia; chronic obstructive airways disease; cardiac arrhythmia; xerostomia; diabetes mellitus; Sjogren's syndrome; Paget's disease; hereditary hematochromatosis; or a non-CNS disorder resulting from normal aging.
- 15 20 76. The method of any of claims 45, 47, 48, 49, 56, 58, 60, 63, 65, or 68, wherein the subject suffers from at least one of a neurologic disorder and a psychiatric disorder.
- 25 77. The method of claim 76, wherein the subject suffers from at least one of Alzheimer's disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, an affective disorder, an anxiety disorder, schizophrenia, cell damage, nerve damage, a CNS infection, a tumor of the brain, a tumor of the spinal cord, a stroke in the brain, a stroke in the spinal cord, a prion disease, a CNS disorder


resulting from ordinary aging, a brain injury, a spinal cord injury, or a combination thereof.

78. The method of any of claims 45, 47, 48, 49, 56, 58, 60, 63, 65, or 68, further comprising administering heme oxygenase, a vector encoding a heme oxygenase, heme oxygenase-1, a vector encoding a heme oxygenase-1, heme oxygenase-2, a vector encoding a heme oxygenase-2, a biliverdin reductase, a vector encoding a biliverdin reductase, a catalase, a vector encoding a catalase, a peroxidase, a vector encoding a peroxidase, or a combination thereof.
79. The method of any of claims 45, 47, 48, 49, 56, 58, 60, 63, 65, or 68, further comprising administering a heme binding protein.
80. The method of claim 79, wherein the heme binding protein comprises hemopexin, a lipoprotein, or a combination thereof.


**Pyrophosphate Protects the mAChR from
Inactivation by the LMW Inhibitor
(n = 3)**




**Pyrophosphate Protects the mACHR
from Inactivation by 2 μ M Heme and 100 μ M Peroxide
(n = 10)**




**Pyrophosphate protects the mAChR from
inactivation by the LMW Inhibitor in a
2-step agonist binding assay.
(n = 4)**




Imidodiphosphate protects the mAChR
from inactivation by the LMW inhibitor
(n = 5)




**Guanylimidodiphosphate Protects the mAChR
from Inactivation by the LMW Inhibitor
(n = 4)**




**Adenylylimidodiphosphate protects the mAChR from
inactivation by the LMW Inhibitor
(n = 3)**




**Tripolyphosphate Protects the mAChR from LMW
Inhibitor inactivation in Atagonist Binding Studies
(n = 4)**




**Bilirubin Protects the mAChR from
Inactivation by the LMW Inhibitor
(n = 4)**




**Bilirubin Protects the mAChR from
Inactivation by 2.0 μ M Heme and 100 μ M Peroxide
(n = 4)**




**Bilirubin Protects the mAChR from LMW Inhibitor
inactivation in the two step agonist (oxotremorine-m)
binding assay
(n = 3)**



**Billiverdin Protects the mAChR
from Inactivation by the LMW Inhibitor
(n = 7)**




**1 μ M Carnosol completely protects the mAChR
from inactivation by the LMW Inhibitor.**




Percentages Represent: Percent Inhibition/Percent Protection


**Quercetin protects the mAChR
from inactivation by the LMW inhibitor
(n = 3)**



**Myricetin Protects the mAChR
from Inactivation by Heme and Peroxide**




Catalase Protects the mAChR from Inactivation by the LMW Inhibitor




15/20

**Catalase Protects the mACHR from
Inactivation by 2.0 μ M Heme and 100 μ M Peroxide**




Glutathione peroxidase protects the mAChR from inactivation by the LMW inhibitor




*Glutathione peroxidase concentration in the presence of the LMW Inhibitor
Percentages represent: Percent Inhibition / Percent Protection

**Glutathione peroxidase protects the mAChR from inactivation
by 2 μ M Heme and 100 μ M H_2O_2**




***Glutathione peroxidase concentration in the presence of Heme and Peroxide
Percentages represent: Percent Inhibition / Percent Protection**

**Pamidronate (the active component of Aredia®)
Protects the mAChR from Inactivation
by the LMW Inhibitor
(n=4)**



*Aredia is a drug manufactured by Novartis and is a bone-resorption inhibitor for treatment of hypercalcaemia. All assay tubes contained 7.94 mM Mannitol a component of Aredia which had no effect on binding or inhibition.

Pyrophosphate Protects the mACHR
from damage by 160 μM PbCl_2



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/082932 A3

(51) International Patent Classification⁷: A61K 31/661, 31/6615, 31/662, 31/7084, 31/706, 31/7076, A61P 33/00

(21) International Application Number: PCT/US01/13931

(22) International Filing Date: 30 April 2001 (30.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/200,843 1 May 2000 (01.05.2000) US
60/230,263 6 September 2000 (06.09.2000) US
60/233,025 15 September 2000 (15.09.2000) US

(71) Applicant: HEALTHPARTNERS RESEARCH FOUNDATION [US/US]: 8100 34th Avenue South, Minneapolis, MN 55440-1524 (US).

(72) Inventors: FREY, William, H., II.: 4800 Centerville Road, Apt. 216, White Bear Lake, MN 55127 (US). FAWCETT, John, Randall; 928 Grand Avenue, # 305, St. Paul, MN 55105 (US). THORNE, Robert, Gary; 2610 Fremont Avenue South #106, Minneapolis, MN 55408 (US). CHEN, Xueqing; 1166 Fifield Avenue, Apt. S2, St. Paul, MN 55108 (US).

(74) Agent: FORREST, Peter, R.; Gray, Plant, Mooty, Mooty & Bennett, P.A., P.O. Box 2906, Minneapolis, MN 55402-2906 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
18 July 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND COMPOSITIONS FOR ENHANCING CELLULAR FUNCTION THROUGH PROTECTION OF TISSUE COMPONENTS

(57) Abstract: Methods and compositions for enhancing cellular function through protection of tissue components, such as receptors, proteins, lipids, nucleic acids, carbohydrates, hormones, vitamins, and cofactors, by administering pyrophosphate analogs or related compounds.

WO 01/082932 A3

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/661 A61K31/6615 A61K31/662 A61K31/7084 A61K31/706
A61K31/7076 A61P33/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Section Ch, Week 198701 Derwent Publications Ltd., London, GB; Class A97, AN 1987-003703 XP002189397 & JP 61 263902 A (RIKAGAKU KENKYUSHO), 21 November 1986 (1986-11-21) abstract</p> <p>-----</p>	16-18

☐ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

6 February 2002

Date of mailing of the international search report

08 MAY 2002

Name and mailing address of the ISA

European Patent Office, P.B: 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herrera, S

INTERNATIONAL SEARCH REPORT

In: tional application No.
PCT/US 01/13931

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-15, 19-59, 62-80
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

16-18

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Continuation of Box I.1

Although claims are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Claims Nos.: 1-15,19-59,62-80

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

It is further pointed out that the subject-matter of claim 16 has only been searched as far as the methods relates to the treatment of plants.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 16-18

Use of pyrophosphate and derivatives thereof for treating bacterial, fungal, algae and algo infections in plants

2. Claims: 60-61

Use of bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavinoid or combinations thereof for treating bacterial, fungal, algae and algo infections in plants

Information on patent family members

PC1, US 01/13931

Form PCT/ISA/210 (patent family annex) (July 1992)